



Localization of classical swine fever virus (CSFV) in the brain of pigs infected with strain Kozlov

Hansen, Mette Sif; Bille-Hansen, Vivi; Uttenthal, Åse; Lohse, Louise; Nielsen, Jens

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Hansen, M. S., Bille-Hansen, V., Uttenthal, Å., Lohse, L., & Nielsen, J. (2011). *Localization of classical swine fever virus (CSFV) in the brain of pigs infected with strain Kozlov*. Abstract from 8th ESVV International Pestivirus Symposium, Hanover, Lower Saxony, Germany. <http://www.pestivirus2011.de/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Localization of classical swine fever virus (CSFV) in the brain of pigs infected with strain Kozlov

Mette Sif Hansen¹, Vivi Bille-Hansen², Åse Uttenthal¹,
Louise Lohse¹, Jens Nielsen¹

¹ Division of Virology, DTU National Veterinary Institute, Lindholm, 4771 Kalvehave, Denmark

² DanaLab, Agern Alle 3, DK-2970 Hørsholm, Denmark

While, some CSFV strains induce moderate to severe affection of the central nervous system (CNS), others do not¹.

The aim of the present experiment was to study putative correlations between the intensity of brain lesions or CSFV-infected cells, – and CNS symptoms, viral RNA load or the number of CD3-positive cells in brain tissue. Nine pigs were inoculated with CSFV Kozlov, 6 pigs were controls. Daily clinical examinations, including CNS scoring, were carried out. The pigs were euthanized on PID 6-10. Necropsy was performed and formalin-fixed brain tissue was examined for CSFV and CD3² by immunohistochemistry.

CNS symptoms in infected pigs were ataxia (n=8), paresis (n=1) and convulsions (n=5).

All infected pigs had brain lesions, like congestion, meningitis with mononuclear cells and/or accumulations of eosinophils, and perivascular cuffings.

All infected pigs had CSFV-infected cells in the brain. The cells were mainly localized around and in vessels in all parts of the cerebrum and cerebellum. The infected cells were endothelium and lymphocytes, however, dendrite-like cells were also seen. Quantitative-RT-PCR³ showed that the viral RNA load of the brain correlated well to that of serum.

CD3-positive cells were seen in infected pigs and controls, but in higher numbers in infected pigs, which also had considerably more CSFV-infected cells than CD3-positive cells.

All inoculated pigs had non-suppurative meningo-encephalitis. The intensity of brain lesions correlated well to the number of CSFV-infected cells. There was no obvious association between the number of lesions or CSFV-infected cells, and – CNS symptoms, the number of CD3-positive cells or viral RNA load in the brain.

References

¹Belak et al., (2008). Acta Vet. Scand., 50, 34

²Modified after Tingstedt et al., (2003). Vet. Immunol. Immunopathol., 94, 123-132

³Rasmussen et al., (2007). J. Gen. Virol., 88, 481-486